

TURN FORM OVER

Food Allergy Action Plan

Name:			D.O.B.:	:	1	Place Student's	
Allergy to:	Allergy to: Picture						
Weight:	ibs. Asthma: □ Y	es (higher risk fo	r a severe reacti	ion) 🗆	l No	Here	
Extremely reactive to the following foods: THEREFORE: If checked, give epinephrine immediately for ANY symptoms if the allergen was likely eaten. If checked, give epinephrine immediately if the allergen was definitely eaten, even if no symptoms are noted.							
ingestion: One or more LUNG: HEART: THROAT: MOUTH: SKIN:	confused Tight, hoarse, trouble bi	e, repetitive cough oulse, dizzy, reathing/swallowingue and/or lips) ferent body areas	ng S	2. C. 3. Be be 4. G. *Antihil are nor severe	AMEDIATI all 911 egin monito elow) ive addition Antihistamii Inhaler (bro asthma istamines & in t to be depend	oring (see box nal medications:*	
MILD SYMPT MOUTH: SKIN: GUT:	Itchy mouth A few hives around mou			2. S h p 3. If	itay with stu ealthcare p arent symptoms bove), USE	IISTAMINE Ident; alert rofessionals and progress (see EPINEPHRINE oring (see box	
Medications/Doses below)							
Epinephrine (brand and dose):							
Antihistamine (brand and dose):							
Other (e.g., inhaler-bronchodilator if asthmatic):							
Monitoring Stay with student; alert healthcare professionals and parent. Tell rescue squad epinephrine was given; request an ambulance with epinephrine. Note time when epinephrine was administered. A second dose of epinephrine can be given 5 minutes or more after the first if symptoms persist or recur. For a severe reaction, consider keeping student lying on back with legs raised. Treat student even if parents cannot be reached. See back/attached for auto-injection technique.							
Parent/Guardian	Signature	Date	Physician/Health	care Pro	vider Signat	ure Date	


Form provided courtesy of FAAN (www.foodallergy.org) 7/2010

EPIPEN Auto-Injector and EPIPEN Jr Auto-Injector Directions

- First, remove the EPIPEN Auto-Injector from the plastic carrying case
- Pull off the blue safety release cap



 Hold orange tip near outer thigh (always apply to thigh)



 Swing and firmly push orange tip against outer thigh. Hold on thigh for approximately 10 seconds.
 Remove the EPIPEN Auto-Injector and massage the area for 10 more seconds



BOY' and the Gey logs, Epihen', Epihen 2-Hak', and Epihen ir Z-Pak' are regulated transments of Dey Pharma, 1.9.

Twinject® 0.3 mg and Twinject® 0.15 mg Directions



Remove caps labeled "1" and "2."

Place rounded tip against outer thigh, press down hard until needle penetrates. Hold for 10 seconds, then remove.



SECOND DOSE ADMINISTRATION: If symptoms don't improve after 10 minutes, administer second dose:

Unscrew rounded tip. Pull syringe from barrel by holding blue collar at needle base.



Put needle into thigh through skin, push plunger down all the way, and remove.



Adrenaclick™ 0.3 mg and Adrenaclick™ 0.15 mg Directions



Remove GREY caps labeled "1" and "2."

 A food allergy response kit should contain at least two doses of epinephrine, other medications as noted by the student's physician, and a copy of this Food Allergy Action Plan.

A kit must accompany the student if he/she is off school grounds (i.e., field trip).

Contacts	
Call 911 (Rescue squad: ()	Phone: ()
Parent/Guardian:	Phone: ()
Other Emergency Contacts Name/Relationship:	Phone: () -
Name/Relationship:	Phone: () -